

## SUMMARY

### A phytosociological synthesis of Mopaneveld

by

Frances du Plessis

Supervisor: Prof. Dr. G.J. Bredenkamp

Submitted in partial fulfilment of the requirements for the degree

### MAGISTER SCIENTIA (Botany)

One of the most critical shortcomings with regard to vegetation utilisation and conservation in southern Africa, is the lack of in-depth knowledge of the ecology of the various vegetation types. Mopaneveld is one of the largest savannas of Africa. However, knowledge of this extensive vegetation type is still limited to local-scale studies within the different *Colophospermum mopane*-hosting countries. The major aim of this study was to analyse existing phytosociological data from Mopaneveld over its entire distribution range. This holistic view of Mopaneveld vegetation contributes to the better understanding of large-scale vegetation continua, which contains identifiable plant communities along environmental gradients. Furthermore, provides a better understanding of ecological processes within this vegetation type, which occupies vast areas over a considerable variation in environmental parameters.

Compatible vegetation data were collected and captured in a TURBOVEG database. Known methods to analyse large vegetation data sets were considered. These methods could however not be applied successfully to the synthesis of the Mopaneveld because all suitable data were not processed to describe plant communities yet. A new method to treat large vegetation data sets was proposed. This method is based on the synthesis of raw data material by means of basic phytosociological principles. The complete data set was analysed by TWINSpan procedures in MEGATAB. Results are presented in synoptic tables. Although the synoptic presentation limits the number of relevés, the number of species remains high. A reduced synoptic table was constructed to accommodate the need for a clear presentation of results in a reduced format. The

method was critically evaluated after which was concluded that the method is only a step in the direction of new methodology in meta-analysis.

A literature review on *Colophospermum mopane* was prepared in order to provide sufficient information on this plant species, which dominates the woody strata of the Mopaneveld through most of its distribution range. Lists of known described plant communities and vegetation types within the Mopaneveld are provided in two appendices. Mopaneveld occurs in eight African countries, which include Angola, Namibia, Botswana, Zambia, Zimbabwe, Malawi, Mozambique and South Africa. A summary on the Mopaneveld in these hosting countries is provided to contribute to the knowledge of Mopaneveld vegetation of its total distribution range.

TWINSPAN classification and subsequent refinement by procedures of the new approach, resulted in the identification of seven vegetation types and six major plant communities in the Mopaneveld of South Africa, Zimbabwe and Namibia. Due to special interest in the Mopaneveld of the South African Lowveld, a synthesis of this vegetation type was prepared separately. Four major plant communities were identified from this procedure of which two are described in this dissertation.

This synthesis revealed interesting notes on the dynamics of Mopaneveld vegetation. Results from the synthesis provided information on the probability that the Mopaneveld is an event-driven system and its dynamics can be explained by non-equilibrial models of vegetation change.

## OPSOMMING

**‘n Fitososiologiese sintese van Mopaneveld**

deur

**Frances du Plessis**

Studieleier: Prof. Dr. G.J. Bredenkamp

Voorgelê ter gedeeltelike vervulling van die vereistes vir die graad

### **MAGISTER SCIENTIA (Plantkunde)**

Een van die grootste tekortkominge in bewaring en grondgebruik in suidelike Afrika is die gebrek aan in-diepte kennis van die ekologie van verskillende plantegroeitipes. Mopaneveld is een van die grootste savannas in Afrika. Kennis van hierdie uitgestrekte plantegroeitipe is nietemin beperk tot plaaslike studies binne die verskillende lande waarin *Colophospermum mopane* voorkom. Die primêre doel van hierdie studie was om bestaande fitososiologiese data van Mopaneveld oor die totale verspreiding daarvan te analiseer. Hierdie holistiese benadering van Mopaneveld plantegroei dra by tot ‘n beter oorsig van grootskaalse plantegroeykontinua wat identifiseerbare plantgemeenskappe bevat langs omgewingsgradiënte. Verder voorsien hierdie studie ook in ‘n beter oorsig van ekologiese prosesse binne die Mopaneveld, wat ‘n groot area beset oor ‘n merkwaardige variasie in omgewingsparameters.

Bruikbare plantegroeydata was versamel en in ‘n TURBOVEG databasis gestoor. Bekende metodes was oorweeg vir die analise van die groot datastel. Hierdie metodes kon egter nie suksesvol toegepas word in die sintese van die Mopaneveld nie aangesien nie alle data geprosesseer en die plantgemeenskappe daarvan opgeskryf is nie. ‘n Nuwe metode oor die hantering van ‘n groot datastel is daarom voorgestel. Die metode is gebaseer op die sintese van roudatamateriaal volgens basiese fitososiologiese beginsels. Die totale datastel was geanaliseer in MEGATAB deur die gebruik van TWINSPAN prosedures. Resultate word voorgestel in sinoptiese tabelle. Die sinoptiese voorstellings beperk die aantal relevés, nietemin bly die aantal spesies hoog. A verkorte sinoptiese tabel is daarom saamgestel sodat resultate duidelik

voorgestel kan word. Die metode was krities ondersoek waarna beslis is dat die voorgestelde metodiek slegs beskou kan word as die eerste stap in die rigting van nuwe metodiek in meta-analise.

‘n Literatuuroorsig van *Colophospermum mopane* is voorberei vir die weergee van nuttige inligting oor die spesie wat die houtagtige komponent van Mopaneveld domineer. Lyste van plantgemeenskappe en plantegroeitipes binne Mopaneveld wat reeds beskryf is, is voorsien. Mopaneveld kom voor in agt Afrika-lande, wat Angola, Namibië, Botswana, Zambië, Zimbabwe, Malawi, Mosambiek en Suid-Afrika insluit. ‘n Opsomming oor die Mopaneveld van hierdie lande is gegee ter gedeeltelike bydrae tot kennis oor Mopaneveld plantegroei, oor die totale verspreidingsgebied.

TWINSpan klassifikasie en die verfyning daarvan op grond van die nuwe metode, het gelei tot die identifisering van sewe plantegroeitipes en ses hoofplantgemeenskappe in die Mopaneveld van Suid-Afrika, Zimbabwe en Namibië. ‘n Afsonderlike sintese van die Suid-Afrikaanse Laeveld Mopaneveld was voorberei. Vier hoof plantgemeenskappe was identifiseer waarvan twee beskryf is in hierdie verhandeling.

Hierdie sintese het verder interessante opmerkings oor die plantegroeidinamika van Mopaneveld opgelewer. Resultate van die sintese het bygedra tot die versterking van spekulasies dat die Mopaneveld gedryf word deur gebeurtenisse. Die dinamika kan met ander woorde voorgestel word deur nie-ekwilibrium modelle vir plantegroei-verandering.

## ACKNOWLEDGEMENTS

Thanks are due to the following institutions and persons for their valuable contribution in the preparation of this dissertation:

- Prof. George Bredekamp for his advice, interest, time, support and valuable guidance
- Dr. Stephan Hennekens, University of Lancaster, The Netherlands, for his valuable advice on the use of TURBOVEG and MEGATAB
- Prof. Laco Mucina, University of the North, Qwa-Qwa campus, for valuable discussions on phytosociology
- Dr. Ben Strohbach, NBRI, Namibia, for his valuable scientific contribution during a field trip to Namibian Mopaneveld
- Mr. Jonathan Timberlake, Biodiversity Foundation of Africa, Zimbabwe, for exchanging some thoughts on Mopaneveld and making available the distribution map of *Colophospermum mopane*
- Dr. Augustine Chikuni, National Herbarium and Botanic Garden, Malawi, for valuable assistance in Mopaneveld literature
- Prof. Braam van Wyk for valuable discussions on the floristics of Mopaneveld
- Prof. M.C. Laker, Department of Soil Science, University of Pretoria, for valuable discussions on soil types associated with Mopaneveld
- All authors of data sets (listed in Table 2) who made their vegetation data available for this synthesis
- Ms Magda Geringer, Department of Geography, University of Pretoria, for her help with the compilation of the geological map
- My husband, Stefan Siebert, for his scientific advice and moral support during the compilation of this dissertation
- Ms. Claudia Ueckermann for her assistance during field surveys
- Ms Martie Dednam, H.G.W.J. Schweickerdt Herbarium, University of Pretoria, for technical assistance in handling the collected specimens;
- The Foundation for Research Development, the Department of Environmental Affairs and Tourism and the University of Pretoria for financial support.

## REFERENCES

- ACOCKS, J.P.H. 1953. Veld Types of South Africa. *Mem. Bot. Surv. S. Afr.* 28: 1–192.
- ACOCKS, J.P.H. 1988. Veld Types of South Africa. 3rd edn. *Mem. Bot. Surv. S. Afr.* 57: 1–146.
- ANDERSON, G.D. & WALKER, B.H. 1974. Vegetation composition and elephant damage in the Sengwa Wildlife Research Area, Rhodesia. *J. sth. Afr. Wildl. Mgmt. Ass.* 4(1): 15–23.
- ARNOLD, T.H. & DE WET, B.C. 1993. Plants of southern Africa: names and distribution. *Mem. Bot. Surv. S. Afr.* 62: 1–825.
- BARBOSA, L.A. GRANDVAUX. 1970. Carta fitogeográfica de Angola. Luanda, Angola. Instituto de Investigação Científica de Angola.
- BECK, N.D. 1998. A wildlife management plan involving ecological studies of plant communities and animal populations for Foskor mine, Phalaborwa. B.Sc. (Hons.) dissertation, University of Pretoria, Pretoria.
- BECKER, T. & JÜRGENS, N. 2000. Vegetation along climate gradients in Kaokoland, North-West Namibia. *Phytocoenologia* 30(3–4): 543–565.
- BEHNKE, R.H. & SCOONS, I. 1993. Rethinking Range Ecology: Implications for rangeland management in Africa. In: Range Ecology at disequilibrium: New models of natural variability and pastoral adaptation in African Savannas, ed. R.H. Behnke (Jr.), I. Scoones & C. Kerven, Ch. 1, pp. 1–30. Overseas Development Institute, London.
- BEHR, C.M. & BREDENKAMP, G.J. 1988. A phytosociological classification of the Witwatersrand National Botanic Garden. *S. Afr. J. Bot.* 54(6): 525–533.
- BEN-SHAHAR, R. 1993. Patterns of elephant damage to vegetation in Northern Botswana. *Biol. Cons.* 65(2): 249–256.
- BEN-SHAHAR, R. 1996. Do elephants over-utilise mopane woodlands in northern Botswana? *Journal of Tropical Ecology* 12: 505–515.
- BEZUIDENHOUT, H. 1996. The major vegetation communities of the Augrabies Falls National Park, Northern Cape. 1. The southern section. *Koedoe* 39(2): 7–24.
- BHIMA, R. & BREDENKAMP, G.J. 1999. The effect of fire on regeneration of *Colophospermum mopane* and *Dalbergia melanoxylon* and on elephant browsing. *Phytocoenologia* 29(4): 469–484.
- BHIMA, R. & BREDENKAMP, G.J. in press. Impact of elephant on *Colophospermum mopane* trees in Liwonde National Park, Malawi. Submitted to *Phytocoenologia*.
- BIGGS, R.C. 1979. The ecology of Chief's Island and the adjacent floodplains of the Okavango Delta, Botswana. M.Sc. dissertation, University of Pretoria, Pretoria.
- BONYONGO, M.C. 1999. Vegetation ecology of the seasonal floodplains in the Okavango Delta, Botswana. M.Sc. thesis, University of Pretoria, Pretoria.



- BOUGHEY, A.S. 1961. The vegetation types of Southern Rhodesia: a reassessment. *Proc. & Trans. of Rhodesia Scientific Assoc.* 49: 54–98.
- BOURLIÈRE, F. 1983. *Ecosystems of the world* Vol. 13, Tropical savannas. Elsevier, Amsterdam, Oxford & New York.
- BREDENKAMP, G.J. & THERON, G.K. 1990. The vegetation of the fersiallitic soils of the Manyeleti Game Reserve. *Coenoses* 5(3): 167–175.
- BREDENKAMP, G.J. & THERON, G.K. 1991. The *Euceo divinori - Acacietum nigricentis*, a new association from the calcareous bottomland clays of the Manyeleti Game Reserve, Eastern Transvaal Lowveld, Gazankulu, South Africa. *Vegetatio* 93: 119–130.
- BREDENKAMP, G.J., DEUTSCHLÄNDER, M.S. & THERON, G.K. 1993. A phytosociological analysis of the *Albizia harveyi - Eucleetum divinori* from sodic bottomland clay soils of the Manyeleti Game Reserve, Gazankulu, South Africa. *S.Afr.J.Bot.* 59(1): 57–64.
- BREDENKAMP, G.J. & DEUTSCHLÄNDER, M.S. 1994. The *Themeda triandrae - Setariaeum incrassatae*, a new association from gabbro in the Manyeleti Game Reserve, Gazankulu, South Africa. *Koedoe* 37(2): 43–57.
- BREDENKAMP, G.J. & BEZUIDENHOUT, H. 1995. A proposed procedure for the analysis of large phytosociological data sets in the classification of South African grasslands. *Koedoe* 38(1): 33–39.
- BREDENKAMP, G.J. & DEUTSCHLÄNDER, M.S. 1995. New azonal syntaxa from the hills and river banks of the Manyeleti Game Reserve, Northern Transvaal Province, South Africa. *Koedoe* 38(1): 41–58.
- BRETELER, F.J., FERGUSON, I.K. GASSON, P.E. & TER WELLE, B.J.H. 1997. *Colophospermum* reduced to *Hardwickia* (Leguminosae-Caesalpinioideae). *Adansonia* 19: 279–291.
- BROPHY, J.J., BOLAND, D.J., VAN DER LINGEN, S. 1992. Essential oils in the leaf, bark and seed of mopane (*Colophospermum mopane*). *South African Forestry Journal* 161: 23–25.
- BROWN, L.R., BREDENKAMP, G.J. & VAN ROOYEN, N. 1995a. The phytosociology of the southern section of Borakalalo Nature Reserve. *Koedoe* 37(2): 59–72.
- BROWN, L.R., BREDENKAMP, G.J. & VAN ROOYEN, N. 1995b. The phytosociology of the western section of Borakalalo Nature Reserve. *Koedoe* 38(2): 49–64.
- BROWN, L.R., BREDENKAMP, G.J. & VAN ROOYEN, N. 1996. The phytosociology of the northern section of Borakalalo Nature Reserve. *Koedoe* 39(1): 9–24.
- BROWN, L.R. 1997. A plant ecological study and wildlife management plan of the Borakalalo Nature Reserve, North-West Province. Ph.D thesis, University of Pretoria, Pretoria.
- BRUELHEIDE, H. 1995. Die Grünlandgesellschaften des Harzes und ihre Standortsbedingungen. Mit einem Beitrag zum Gliederungsprinzip auf der Basis von statistisch ermittelten Artengruppen. *Diss. Bot.* 244: 1–338.

- BRUELHEIDE, H. 2000. A new measure of fidelity and its application to defining species groups. *J. Veg. Sci.* 11: 167–178.
- CAMPBELL, B.M. & DU TOIT, R.F. 1994. Vegetation patterns and the influence of small-scale farmers in a semi-arid savanna area in Zimbabwe. *Kirkia* 15(1): 10–32.
- CARTER, R.N. & PRINCE, S.D. 1988. Distribution limits from a demographic viewpoint. In: Plant Population Ecology, ed. A.J. Davy, M.J. Hutchings & A.R. Warkinson. *British Ecological Society Symposium* 28: 165–184.
- CHESSON, P.L. & CASE, T.J. 1986. Overview: non-equilibrium community theories: chance, variability, history and coexistence. In: Community Ecology, ed. J. Diamond & T.J. Case, Ch. 13. Harper & Row Publishers, New York.
- CHIKUNI, A.C. 1996. Conservation status of mopane woodlands in Malawi: a case study of Mua-Tsanya Forest Reserve. In: The Biodiversity of African Plants, eds. L.J.G van der Maesen, *et al.* pp.250–258. Kluwer Academic Publishers.
- CHOINSKI, J.S. & TUOHY, J.M. 1991. Effect of water potential and temperature on the germination of four species of African savanna trees. *Ann. Bot.* 68: 227–233.
- CHYTRÝ, M., TICHÝ, L. & HOLT, J. in press. On the fidelity, synoptic tables, and diagnostic species in phytosociology. Submitted to *J. Veg. Sci.*
- CLEMENTS, F.E. 1916. Plant succession: an analysis of the development of vegetation. *Washington: Carnegie Institute Pub.* 242: 1–512.
- COE, K.H. 1991. Effects of thinning on *Colophospermum mopane* in an indigenous woodland setting. *Journal of the Forestry Association of Botswana* 1991: 47–57.
- COE, K.H. 1992. Agroforestry alternatives. Managing natural woodlands in dryland Botswana. *Agroforestry Today* 1992: 6–8.
- COETZEE, B.J. 1983. Phytosociology, vegetation structure and landscapes of the central district, Kruger National Park, South Africa. *Dissertationes Botanicae* 69: 1–456.
- COLE, M.M. 1986. The Savannas: biogeography and geobotany. Academic Press, London.
- COOK, E. 1996. Implications of modern successional theory for habitat typing: a review. *Forest Science* 42(1): 67–75.
- CORREIA, R.I. De S. 1976. The main vegetation types of Kaokoland, Northern Damaraland and a description of some transects in Owamboland, Etosha and North-western South West Africa. Unpublished report.
- COWLING, R.M. & HILTON-TAYLOR, C. 1994. Patterns of plant diversity and endemism in southern Africa: an overview. *Strelitzia* 1: 31–52.
- DAVIS, S.D., HEYWOOD, V.H. & HAMILTON, A.C. 1994. Centres of plant diversity. A guide and strategy for their conservation. Volume 1. IUCN Publications Unit, Cambridge.



- DEANGELIS, D.L. & WATERHOUSE, J.C. 1987. Equilibrium and nonequilibrium concepts in ecological models. *Ecological Monographs* 57(1): 1–21.
- DEKKER, B. & VAN ROOYEN, N. 1995. The physical environment and plant communities of the Messina Experimental Farm. *S. Afr. J. Bot.* 61(3): 158–167.
- DEKKER, B. & SMIT, G.N. 1996. Browse production and leaf phenology of some trees and shrubs in different *Colophospermum mopane* savanna communities. *African Journal of Range & Forage Science* 13(1): 15–23.
- DEKKER, B., VAN ROOYEN, N. & BOTHMA, J. DU P. 1996. Habitat partitioning by ungulates on a game ranch in the Mopani veld. *S. Afr. J. Wildl. Res.* 26(4): 117–122.
- DE LA HUNT, T.E. 1954. The value of browse shrubs and bushes in the lowveld of the Gwanda area, South Rhodesia. *Rhodesian Agricultural Journal* 51: 251–262.
- DODD, J.L. 1994. Desertification and degradation in Sub-Saharan Africa: The role of livestock. *Bioscience* 44(1): 28–34.
- DONALDSON, C.H. 1979. Goats and/or cattle on Mopani veld. *Proc. Grassl. Soc. Sth. Afr.* 14: 119–123.
- DONALDSON, C.H., ROOTMAN, G.T. & VAN DER MERWE, P. 1984. Ekologie van mopanieveld: sekondêre suksessie op ontblote mopanieveld. Department of Agriculture Final Report. Research Project T5411/27/1/1, Transvaal Region, Pretoria.
- DUDLEY, C.O. 1994. The flora of Liwonde National Park, Malawi. *Proc. XIIIth Plenary Meeting AETFAT, Malawi* 2: 1485–1509.
- DU PLESSIS, W.P., BREDEKAMP, G.J. & TROLLOPE, W.S.W. 1998. Response of herbaceous species to a degradation gradient in the western region of Etosha National Park, Namibia. *Koedoe* 41(1): 9–18.
- DU PLESSIS, F. 2000. The need for compatible floristic data in comprehensive syntaxonomical synthesis with special reference to Mopaneveld in Africa. Unpublished presentation, 26<sup>th</sup> South African Association of Botanists (SAAB) conference, Potchefstroom.
- DU TOIT, R. 1993. Reconnaissance vegetation survey of the Chewore – Angwa – Kanyemba area of the Zambezi Valley, Zimbabwe. *Kirkia* 14(1): 61–77.
- ECKHARDT, H.C., VAN ROOYEN, N. & BREDEKAMP, G.J. 1996. The plant communities and species richness of the *Alepidea longifolia* - *Monocymbium cerasiiforme* High-altitude Grassland of northern Kwa-Zulu-Natal. *Koedoe* 39(1): 53–68.
- EGLER, F.E. 1954. Philosophical and practical considerations of the Braun-Blanquet system of phytosociology. *Castanea* 19: 45–60.
- ELLIS, J.E. & SWIFT, D.M. 1988. Stability of African pastoral ecosystems: Alternate paradigms and implications for development. *J. Range Manage.* 41(6): 450–459.

- ERKKILÄ, A. & SIISKONEN, H. 1992. Forestry in Namibia. *Silve Carelica* 20. University of Joensuu, Faculty of Forestry.
- FANSHAWE, D.B. 1969. The vegetation of Zambia. *Forest Research Bulletin* 7: 1–67. Division of Forest Research, Kitwe, Zambia.
- FARRELL, J.A.K. 1968a. Preliminary notes on the vegetation of lower Sabi-Lundi basin, Rhodesia. *Kirkia* 6(2): 233–248.
- FARRELL, J.A.K. 1968b. Preliminary notes on the vegetation of southern Gokwe district, Rhodesia. *Kirkia* 6(2): 249–257.
- FRASER, S.W., VAN ROOYEN, T.H. & VERSTER, E. 1987. Soil-plant relationships in the Central Kruger National Park. *Koedoe* 30: 19–33.
- FRIEDEL, M.H. 1991. Range condition assessment and the concept of thresholds: A viewpoint. *J. Range Manage.* 44(5): 422–426.
- GAUGH, H.G. 1982. *Multivariate analysis in community ecology*. Cambridge University Press, Cambridge.
- GENEVA EXECUTIVE CENTRE. 1994. *Convention on biological diversity*. Châtelaine, Switzerland.
- GERTENBACH, W.P.D. 1983. Landscapes of the Kruger National Park. *Koedoe* 26: 9–121.
- GERTENBACH, W.P.D. 1987. 'n Ekologiese studie van die suidelikste Mopanieveld in die Nasionale Krugerwildtuin. D.Sc. thesis, University of Pretoria, Pretoria.
- GIESS, W. 1971. A preliminary vegetation map (1: 3 000 000, coloured) of South West Africa. *Dinteria* 4: 5–114.
- GIESS, W. 1998. A preliminary vegetation map of Namibia. *Dinteria* 4: 1–112.
- GUY, P.R. 1975. Notes on the vegetation types of the Zambezi Valley, Rhodesia, between the Kariba and Mpata gorges. *Kirkia* 10: 543–557.
- GUY, P.R. 1981. Changes in the biomass and productivity of woodlands in the Sengwe Wildlife Research Area, Zimbabwe. *Journal of Applied Ecology* 18: 507–519
- HENNEKENS, S.M. 1996a. TURBO(VEG): software package for input, processing, and presentation of phytosociological data. User's guide. Version July 1996. IBN-DLO, Wageningen, and Lancaster University, Lancaster.
- HENNEKENS, S.M. 1996b. MEGATAB: a visual editor for phytosociological tables. User's guide. Version October 1996. IBN-DLO, Wageningen, and Lancaster University, Lancaster.
- HENNING, A.C. & WHITE, R.E. 1974. A study of the distribution of *Colophospermum mopane* (Kirk ex Benth.) Kirk ex J. Léonard: the interactions of nitrogen, phosphorous and soil moisture stress. *Proc. Grassl. Soc. SA*. 9: 53–60.
- HILL, M.O. 1979a. DECORANA – A FORTRAN program for detrended correspondence analysis and reciprocal averaging. Cornell University Ithaca, N.Y.

- HILL, M.O. 1979b. TWINSPAN – A FORTRAN program for arranging multivariate data in an ordered two-way table by classification of individuals and attributes. Cornell University Ithaca, N.Y.
- HINES, C. & BURKE, A. 1997. Vegetation survey of NOLIDEP pilot communities. Prepared for the Northern Regions Livestock Development Project, Ministry of Agriculture, Water and Rural development, Republic of Namibia.
- HIN, C.J. 2000. A natural resource inventory of Sango Ranch, Save Valley Conservancy, Zimbabwe. M.Sc. dissertation, University of Pretoria, Pretoria.
- HUNTLEY, B.J. 1974. Outlines of wildlife conservation in Angola. *J. sth. Afr. Wildl. Mgmt. Ass.* 4(3): 157–166.
- ILLIUS, A.W. & O’CONNOR, T.G. 1999. On the relevance of non-equilibrium concepts to arid and semiarid grazing systems. *Ecological Applications* 9(3): 798–813.
- JARMAN, P.J. & THOMAS, P.I. 1969. Observations on the distribution and survival of mopane (*Colophospermum mopane* (Kirk ex Benth.) Kirk ex J. Léonard) seeds. *Kirkia* 7: 103–107.
- JELTSCH, F., MILTON, S.J., DEAN, W.R.J. & VAN ROOYEN, N. 1996. Tree spacing and coexistence in semiarid savannas. *J. Ecol.* 84: 583–595.
- JELTSCH, F., MILTON, S.J., DEAN, W.R.J., VAN ROOYEN, N. & MOLONEY, K.A. 1998. Modelling the impact of small-scale heterogeneities on tree-grass coexistence in semi-arid savannas. *J. Ecol.* 86: 780–793.
- JORDAAN, A. & WESSELS, D.C.J. 1999. The aril of *Colophospermum mopane*. Its role during seed germination and fruit opening. *S. Afr. J. Bot.* 65(5&6): 392–397.
- JORDAAN, A., DU PLESSIS, H.J. & WESSELS, D.C.J. 2000. Roots of *Colophospermum mopane*. Are they infected by rhizobia? *S. Afr. J. Bot.* 66(2): 128–130.
- JOUBERT, E. 1971. The physiographic, edaphic and vegetative characteristics found in the western Etosha National Park. *Madoqua* 1(4): 5–32.
- JOUBERT, S.C.J. 1976. The population ecology of the roan antelope *Hippotragus equinus equinus* (Desmarest) in the Kruger National Park. D.Sc. thesis, University of Pretoria, Pretoria.
- KENNEDY, A.D. 2000. Wildlife reduces elephant herbivory on *Colophospermum mopane* (Fabaceae). *Afr. J. Ecol.* 38: 175–177.
- KENT, M. & COKER, P. 1995. Phytosociology and the Zurich-Montpellier (Braun-Blanquet) school of subjective classification. In: *Vegetation description and analysis: a practical approach*, ed. M. Kent & P. Coker, 2nd edn, Ch. 7, pp. 245–275. John Wiley & Sons, England.
- KLOK, C.J. & CHOWN, S.L. 1999. Assessing the benefits of aggregation – thermal biology and water relations of anomalous emperor moth caterpillars. *Functional Ecology* 13(3): 417–427.
- KNOOP, W.T. & WALKER, B.H. 1985. Interactions of woody and herbaceous vegetation in a southern African savanna. *J. Ecol.* 73: 235–253.

- KRAUSKOPF, K.B. 1967. Introduction to geochemistry. McGraw-Hill Book Company, New York.
- LAYCOCK, W.A. 1991. Stable states and thresholds of range condition on north American rangelands: a viewpoint. *J. Range Manage.* 44(5): 427-433.
- LÉONARD, J. 1949. Notulae Systematicae IV: Caesalpiniaceae-Amherstieae africanae americanaeque. *Bulletin du Jardin Botanique de L'état (Bruxelles)* 19: 388–391.
- LÉONARD, J. 1999. *Colophospermum* n'est pas synonyme d'*Hardwickia* (Caesalpiniaceae). Conclusion d'une méthode objective de travail. *Bill. Jard. Bot. Nat. Belg* 67: 21–43.
- LE ROUX, C.J.G. 1980. Vegetation classification and related studies in the Etosha National Park. D.Sc thesis, University of Pretoria, Pretoria
- LE ROUX, C.J.G., GRUNOW, J.O., MORRIS, J.W., BREDEKAMP, G.J. & SCHEEPERS, J.C. 1988. A classification of the vegetation of the Etosha National Park. *S. Afr. J. Bot.* 54(1): 1–10.
- LE ROUX, A., SMIT, G.N. & SWART, J.S. 1994. Root biomass of a dense stand of *Colophospermum mopane*. *Bull. Grassld. Soc. Sth. Afr.* 5(1): 50–51
- LEWIS, D.M. 1987. Elephant response to early burning in mopane woodland, Zambia. *S. Afr. J. Wildl. Res.* 17(2): 33–40.
- LEWIS, D.M. 1991. Observations of tree growth, woodland structure and elephant damage on *Colophospermum mopane* in Luangwa Valley, Zambia. *Afr. J. Ecol.* 29: 207–221.
- LOCK, J.M. 1989. Legumes of Africa: a check-list. Royal Botanic Gardens, Kew.
- LOUW, A.J. 1970. 'n Ekologiese studie van Mopanie-veld Noord van die Soutpansberg. D.Sc. thesis, University of Pretoria, Pretoria.
- LOW, A.B. & REBELO, A.G. 1998. Vegetation of South Africa, Lesotho and Swaziland: A companion to the Vegetation Map of South Africa, Lesotho and Swaziland. Department of Environmental Affairs & Tourism, Pretoria.
- MADAMS, R.W. 1990. The biogeography of *Colophospermum mopane* (Kirk ex Benth.) Kirk ex J. Léonard at its distribution limit in eastern Botswana. Ph.D. thesis, Queen Mary Westfields College, University of London, London.
- MADZIBANE, J. & POTGIETER, M.J. 1999. Uses of *Colophospermum mopane* (Leguminosae: Caesalpinioideae) by the Vhavenda. *S. Afr. J. Bot.* 65(5&6): 440–443.
- MALAN, J.W. & VAN WYK, A.E. 1993. Bark structure and preferential bark utilisation by the African elephant. *LAWA Journal* 14(2): 173–185.
- MAPAURE, I. 1994. The distribution of *Colophospermum mopane* (Leguminosae – Caesalpinioideae) in Africa. *Kirkia* 15(1): 1–5.
- MENDELSON, J. & ROBERTS, C. 1997. An environmental profile and atlas of Caprivi. Directorate of Environmental Affairs, Namibia.

- MENTIS, M.T., GROSSMAN, D., HARDY, M.B., O'CONNOR, T.G. & O'REAGAN, P.J. 1989. Paradigm shifts in South African range science, management and administration. *S. Afr. J. Science* 85: 684–687.
- MILTON, S.J., DEAN, W.R.J., DU PLESSIS, M.A. & SIEGFRIED, W.R. 1994. A conceptual model of arid rangeland degradation. *BioScience* 44(2): 70–76.
- MOTSHEGWE, S.M., HOLMBACK, J. & YEBOAH, S.O. 1998. General properties and the fatty acid composition of the oil from the mophane caterpillar, *Imbrasia belina*. *Journal of the American Oil Chemists Society*. 75(6): 725–728.
- MUCINA, L. 1997. Classification of vegetation: Past, present and future. *J. Veg. Sci.* 8: 751–760.
- NEL, P.J., BREDENKAMP, G.J. & VAN ROOYEN, N. 1993. Ecological status of grass species in the red turfveld of the Springbok Flats Turf Thornveld, Transvaal. *S. Afr. J. Bot.* 59(1): 45–49.
- O'CONNOR, T.G. 1985. A synthesis of long term experiments concerning the grass layer of southern African savannas. *S. Afr. Nat. Sci. Prog. Rep.* 114: 1–126.
- O'CONNOR, T.G. & CAMPBELL, B.M. 1986. Classification and condition of the vegetation types of the Nyahungwe area on the Lundi River, Gonarezhou National Park, Zimbabwe. *S. Afr. J. Bot.* 52(2): 117–123.
- O'CONNOR, T.G. 1992. Woody vegetation–environment relations in a semi-arid savanna in the northern Transvaal. *S. Afr. J. Bot.* 58(4): 268–274.
- O'CONNOR, T.G. & ROUX, P.W. 1995. Vegetation changes (1949–71) in a semi-arid, grassy dwarf shrubland in the Karoo, South Africa: influence of rainfall variability and grazing by sheep. *Journal of Applied Ecology* 32: 612–626.
- O'CONNOR, T.G. 1999. Impact of sustained drought on a semi-arid *Colophospermum mopane* savanna. *African Journal of Range & Forage Science* 15(3): 83–91.
- OELOFSE, J., BROCKETT, B.H., BIGGS, H.C. & EBERSOHN, C. 2000. The effect of drought and post-fire grazing on the herbaceous layer of shrub-mopane veld on basalt in the Kruger National Park, South Africa. *V<sup>7</sup><sup>th</sup> International Rangeland Congress Proceedings* 1: 505–507.
- OWEN-SMITH, N., COOPER, S.M., NOVELLIE, P.A. 1983. Aspects of the feeding ecology of a browsing ruminant: the kudu. *South African Journal of Animal Science* 13: 35–38.
- PALGRAVE, K.C. 1983. *Trees of Southern Africa*, 5th edn. Struik, Cape Town.
- PALMER, A.R. & VAN STADEN, J.M. 1992. Predicting the distribution of plant communities using annual rainfall and elevation: an example from southern Africa. *J. Veg. Sci.* 3: 261–266.
- POORE, M.E.D. 1956. The use of phytosociological methods in ecological investigations. IV. General discussion on phytosociological problems. *J. Ecol.* 43: 606–651.
- POTGIETER, M. MADZIBANE, J., MASHABANE, L. & WESSELS, D. 2001. Mopane-veld: Can we afford to loose this valuable veld type? *Veld & Flora* 87(2): 78–79.



- PRIOR, J. & CUTTER, D. 1996. Africa's shrinking savannas.  
<http://ekolserv.vo.slu.se/Docs/www/Subject/Agroforestry/>
- RATTRAY, J.M. & WILD, H. 1961. Vegetation map of the Federation of Rhodesia and Nyasaland. *Kirkia* 2: 94–104.
- RATTRAY, J.M. 1962. Vegetation types of Southern Rhodesia. *Kirkia* 2: 68–93.
- RATTRAY, J.M. 1963. Effect of climate on vegetation with particular reference to southern Rhodesia. General File, Government Printer, Salisbury.
- ROGERS, C.M.L. 1993. A woody vegetation survey of Hwange National Park. Unpublished report, Department of National Parks and Wildlife Management, Zimbabwe.
- ROSS, J.H. 1977. *Colophospermum*. *Flora of Southern Africa* 16(2): 16–19.
- ROTHAUGE, A. 2000. New ecological perceptions of arid rangelands. *Agricola* 2000: 49–56.
- SCHAMINEÉ, J.H.J. & STORTELDER, A.H.F. 1996. Recent developments in phytosociology. *Acta Bot. Neerl.* 45(4): 443–459.
- SCHMIDT, A.G., THERON, G.K. & VAN HOVEN, W. 1993. The phytosociology and structure of vegetation near Villa Nora, north-western Transvaal, South Africa. *S.Afr.J.Bot.* 59(5): 500–510.
- SCHOLES, R.J. 1990. The regrowth of *Colophospermum mopane* following clearing. *J. Grassl Soc. Sth. Afr.* 7: 147–151.
- SCHOLES, R.J. & WALKER, B.H. 1993. African savannas: an overview. In: *An African Savanna: synthesis of the Nylsvley study*, eds. R.J. Scholes & B.H. Walker, pp. 2–16. Cambridge University Press.
- SCHOLES, R.J. 1997. Savanna. In: *Vegetation of Southern Africa*, eds. R.M. Cowling, D.M. Richardson & S.M. Pierce. Cambridge University Press.
- SHARMA, B.D., TEWARI, J.C., GUPTA, I.C. & HARSH, L.N. 1989. *Colophospermum mopane*: an exotic tree for the arid-zone. *Indian Farming* 39(6): 5–6.
- SIEBERT, S.J. 1998. Ultramaphic substrates and floristic patterns in Sekhukhuneland, South Africa. M.Sc. dissertation, University of Pretoria.
- SKARPE, C. 1992. Dynamics of savanna ecosystems. *J. Veg. Sci.* 3: 293–300.
- SMIT, C.M.; BREDEKAMP, G.J.; VAN ROOYEN, N. VAN WYK, A.E. & COMBRINCK, J.M. 1997. Vegetation of the Witbank Nature Reserve and its importance for conservation of threatened Rocky Highveld Grassland. *Koedoe* 40(2): 85–104.
- SMIT, G.N. 1994. The influence of intensity of tree thinning on Mopani veld. Ph.D thesis, University of Pretoria, Pretoria.
- SMIT, G.N. & SWART, J.S. 1994. Influence of leguminous and non-leguminous woody plants on the herbaceous layer and soil under varying competition regimes in Mixed Bushveld. *African Journal of Range and Forage Science* 11(1): 27–33.



- SMIT, G.N., SWART, J.S. & LE ROUX, A. 1994. Root biomass, spatial distribution and relations with aboveground leaf biomass of *Colophospermum mopane*. *Bull. Grassl. Soc. Sth. Afr.* 5(1): 32–39.
- SMIT, G.N., RETHMAN, N.F.G. & MOORE, A. 1996. Vegetative growth, reproduction, browse production and response to tree clearing of woody plants in an African savanna. *Afr. J. Range For. Sci.* 13(2): 78–88.
- SMIT, G.N. & RETHMAN, N.F.G. 1998a. Root biomass, depth distribution and relations with leaf biomass of *Colophospermum mopane*. *S. Afr. J. Bot.* 64(1): 38–43.
- SMIT, G.N. & RETHMAN, N.F.G. 1998b. The influence of tree thinning on the reproduction dynamics of *Colophospermum mopane*. *S. Afr. J. Bot.* 64(1): 25–29.
- SMITH, V.R. 1972. A pot culture investigation into the effect of *Colophospermum mopane* growth on the chemical properties of four selected soils. B.Sc. dissertation, University of the Witwatersrand, Johannesburg.
- SMITH, P.P., TIMBERLAKE, J.R. & VAN WYK, A.E. 1998. Proposal to conserve the name *Colophospermum* against *Hardwickia* (Leguminosae, Caesalpinioideae). *Taxon* 47: 751–752.
- SPRUGEL, D.G. 1991. Disturbance, equilibrium, and environmental variability: What is 'natural' vegetation in a changing environment? *Biol. Cons.* 58: 1–18.
- STODDART, L.A., SMITH, A.D. & BOX, T.W (eds). 1975. Range Management. 3rd edn, McGraw-Hill, New York.
- STYLES, C.V. & SKINNER, J.D. 1996. Possible factors contributing to the exclusion of saturniid caterpillars (mopane worms) from a protected area in Botswana. *Afr. J. Ecol.* 34(3): 276–283.
- STYLES, C.V. & SKINNER, J.D. 1997. Seasonal variations in the quality of mopane leaves as a source of browse for mammalian herbivores. *Afr. J. Ecol.* 35: 254–265.
- STYLES, C.V. & SKINNER, J.D. 2000. The influence of large mammalian herbivores on growth form and utilization of mopane trees, *Colophospermum mopane*, in Botswana's Northern Tuli Game Reserve. *Afr. J. Ecol.* 38: 95–101.
- TER BRAAK, C.J.F. 1995. Ordination. In: Data analysis in community and landscape ecology, eds. R.H.G. Jongman, C.J.F. Ter Braak & O.F.R. Van Tongeren, Ch. 5, pp. 91–173, Cambridge University Press.
- THOMPSON, J.G. 1960. A description of the growth habits of mopani in relation to soil and climatic conditions. *Proceedings of the First Federal Science Congress* 1: 181–186. Rhodesia Scientific Association, Salisbury.
- TICHÝ, L. 2001. JUICE – program for vegetation analysis and classification. *Ann. Bot. (Roma)*.
- TIMBERLAKE, J.R. 1980. Vegetation map of South East Botswana. Report of the Division of Land Utilisation, Department of Agricultural Field Services, Ministry of Agriculture, Gaborone, Botswana.

- TIMBERLAKE, J.R. & MAPAURE, I. 1992. Vegetation and its conservation in the eastern mid-Zambezi valley, Zimbabwe. *Transactions of the Zimbabwe Scientific Association* 66: 1–14.
- TIMBERLAKE, J.R., NOBANDA, N. & MAPAURE, I. 1993. Vegetation survey of the communal lands–north and west Zimbabwe. *Kirkia* 14(2): 171–270.
- TIMBERLAKE, J.R. 1995. *Colophospermum mopane*: annotated bibliography and review. *The Zimbabwe Bulletin of Forestry Research* 11: 1–49.
- TIMBERLAKE, J.R. 1996. *Colophospermum mopane* - a tree for all seasons. In: Sustainable Management of Indigenous Forests in the Dry Tropics, eds. P.T. Mushove, E.M. Shumba & F. Matose, pp. 201–210. Zimbabwe Forestry Commission/SAREC, Harare.
- TIMBERLAKE, J.R. 1999. *Colophospermum mopane*: an overview of current knowledge. In: African Plants: Biodiversity, Taxonomy and Uses, eds. J.R. Timberlake & S. Kativu, pp. 565–571. Royal Botanic Gardens, Kew.
- VAN DER MAAREL, E., ESPEJEL, I. & MORENO-CASASOLA, P. 1987. Two-step vegetation analysis based on very large data sets. *Vegetatio* 68: 139–143.
- VAN DER MAAREL, E. 1990. The Journal of Vegetation Science: a journal for all vegetation scientists. *J. Veg. Sci.* 1: 1–4.
- VAN DER MERWE, J.H. 1983. National Atlas of South West Africa (Namibia). Directorate Development Co-ordination, SWA.
- VAN DER MEULEN, F. 1979. Plant sociology of the western Transvaal Bushveld, South Africa. A syntaxonomical and synecological study. *Dissertationes Botanicae* 49: 1–192.
- VAN OUDTSHOORN, F. 1999. Guide to grasses of southern Africa. Briza, Pretoria, South Africa.
- VAN ROOYEN, N. 1978. 'n Ekologiese studie van die plantgemeenskappe van die Punda Milia-Pafuri-Wambiyagebied in die Nasionale Krugerwildtuin. M.Sc. dissertation, University of Pretoria, Pretoria.
- VAN ROOYEN, N., THERON, G.K. & GROBBELAAR, N. 1981a. A floristic description and structural analysis of the plant communities of the Punda Milia-Pafuri-Wambiya area in the Kruger National Park, Republic of South Africa: 1. The Higrphilous communities. *Jl S. Afr. Bot.* 47(2): 213–246.
- VAN ROOYEN, N., THERON, G.K. & GROBBELAAR, N. 1981b. A floristic description and structural analysis of the plant communities of the Punda Milia-Pafuri-Wambiya area in the Kruger National Park, Republic of South Africa: 2. The Sandveld communities. *Jl S. Afr. Bot.* 47(3): 405–449.
- VAN ROOYEN, N., THERON, G.K. & GROBBELAAR, N. 1981c. A floristic description and structural analysis of the plant communities of the Punda Milia-Pafuri-Wambiya area in the Kruger National Park, Republic of South Africa: 3. The *Colophospermum mopane* communities. *Jl S. Afr. Bot.* 47(4): 585–626.

- VAN ROOYEN, N. & BREDEKAMP, G.J. 1998. Savanna Biome. In: Vegetation of South Africa, Lesotho and Swaziland, eds. A.B. Low & A.G. Rebelo. Department of Environmental Affairs & Tourism, Pretoria.
- VAN VOORTHUIZEN, E.G. 1976. The mopane tree. *Botswana Notes and Records* 8: 223–230.
- VAN WYK, B. & VAN WYK, P. 1997. Field guide to trees of southern Africa. Struik, Cape Town.
- VAN WYK, B-E. & GERICKE, N. 2000. People's Plants. A guide to useful plants of southern Africa. Briza Publishers, Pretoria.
- VENTER, F.J. & GERTENBACH, W.P.D. 1986. A cursory review of the climate and vegetation of the Kruger National Park. *Koedoe* 29: 139–148.
- VILJOEN, P.J. 1980. Veldtipes, verspreiding van die groter soogdiere, en enkele aspekte van die Ekologie van Kaokoland. M.Sc. dissertation, University of Pretoria, Pretoria.
- VISSER, N., VAN HOVEN, W. & THERON, G.K. 1996. The vegetation and identification of management units of the Honnet Nature Reserve, Northern Province, South Africa. *Koedoe* 39(1): 25–42.
- WALKER, B.H., MATTHEWS, D.A. & DYE, P.J. 1986. Management of grazing systems—existing versus an event-orientated approach. *S. Afr. J. Sc.* 82: 112–113.
- WEARE, P.R. & YALALA, A. 1971. Provisional vegetation map of Botswana. *Botswana Notes and Records* 3: 131–147.
- WERGER, M.J.A. 1974. On concepts and techniques applied in the Zürich-Montpellier method of vegetation survey. *Bothalia* 11(3): 309–323.
- WERGER, M.J.A. & COETZEE, B.J. 1978. The Sudano-Zambezian Region. In: Biogeography and ecology of southern Africa, ed. M.J.A. Werger, Ch. 10, pp. 301–462. Junk, The Hague.
- WESTFALL, R.H., VAN ROOYEN, N. & THERON, G.K. 1985. The plant ecology of the farm Groothoek, Thabazimbi District. *Bothalia* 15(3&4): 655–688.
- WESTHOFF, V. & VAN DER MAAREL, E. 1982. The Braun-Blanquet approach. In: Classification of plant communities, ed. R.H. Whittaker, The Hague: Junk.
- WESTOBY, M.; WALKER, B. & NOY-MEIR, I. 1989. Opportunistic management for rangelands not at equilibrium. *J. Range Manag.* 42(4): 266–274.
- WESTFALL, R.H., VAN ROOYEN, N. & THERON, G.K. 1985. The plant ecology of the farm Groothoek, Thabazimbi District. II. Classification. *Bothalia* 15(3&4): 655–688.
- WESTOBY, M.; WALKER, B. & NOY-MEIR, I. 1989. Opportunistic management for rangelands not at equilibrium. *J. Range Manag.* 42(4): 266–274.
- WHITE, F. 1983. The vegetation of Africa, a descriptive memoir to accompany the UNESCO/AETFAT/UNSO vegetation map of Africa. *Natural Resources Research* 20: 1–356.
- WHITTAKER, R.H. 1962. Classification of natural communities. *Bot. Rev.* 28: 1–239.

- WHITTAKER, R.H. 1980. Classification of plant communities, 2nd edn. Dr W. Junk, London.
- WIGGINS, D.A. 1997. Fluctuating asymmetry in *Colophospermum mopane* leaves and oviposition preference in an African silk moth *Imbrasia belina*. *Oikos* 79(3): 484–488.
- WILD, H. & BARBOSA, L.A. GRANDVAUX. 1967. Vegetation map of the Flora Zambesiaca region. *Flora Zambesiaca* supplement. Harare, Zimbabwe.
- WINTERBACH, R. 1998. A phytosociological synthesis of *Acacia tortilis* communities in the North-western savanna of South Africa. M.Sc. dissertation, University of Pretoria, Pretoria, South Africa.
- WOODWARD, F.I. 1986. Climate and vegetation. In: Climate and plant distribution. Cambridge University Press, Cambridge.
- WITKOWSKI, E.T.F. & O'CONNOR, T.G. 1996. Topo-edaphic, floristic and physiognomic gradients of woody plants in a semi-arid African savanna woodland. *Vegetatio* 124: 9–23.